



# Compound Formulation Active Pharmaceutical Ingredients and the Associated Adverse Effects and Potential Contraindications/Warnings and Precautions

## Moxifloxacin + Bromfenac

### Moxifloxacin HCl

#### Possible Adverse Reaction

Conjunctivitis, Decreased visual acuity, Dry eye, Keratitis, Ocular discomfort, Ocular hyperemia, Ocular pain, Ocular pruritus, Subconjunctival hemorrhage, and Tearing, Anaphylaxis, Growth of resistant organisms, Fever, Increased cough, Infection, Otitis media, Pharyngitis, Rash and Rhinitis.

#### Potential Contraindications / Warnings and Precautions

Patients less than 1 year, Lactation, Pregnancy. Known hypersensitivity to flouroquinolones or any of the components. Growth of resistant organisms with prolonged use.

### Bromfenac

#### Possible Adverse Reaction

Anterior chamber eye inflammation, Headache, Vitrious floaters, Iritis, Eye pain, Ocular hypertension, Burning sensation, Conjunctival hyperemia, Corneal erosion, Corneal perforation, Corneal thinning, Corneal ulcer, Epithelial keratopathy, Eye irritation, Eye pruritus, Eye redness, Hypersensitivity reaction, Keratitis, Prolonged bleeding, Stinging sensation, Abnormal sensation in eyes, Anaphylaxis

#### Potential Contraindications / Warnings and Precautions

Lactation, Pediatric patients, Pregnancy, Slow or Delayed Healing (NSAIDS), Potential for cross-sensitivity (NSAIDS) or any of the components. Increased bleeding time of ocular tissue, Keratitis, Corneal reactions. Corneal adverse events are increased in patients with dry eye syndrome, rheumatoid arthritis, repeat ocular surgeries in a short time, and diabetes mellitus.