



Compound Formulation Active Pharmaceutical Ingredients and the Associated Adverse Effects and Potential Contraindications/Warnings and Precautions

Dexamethasone Phosphate + Moxifloxacin HCl

Dexamethasone Sodium Phosphate

Possible Adverse Reaction

Ocular discomfort, Increased IOP, Conjunctival hemorrhage, Conjunctival hyperemia, Anterior chamber inflammation, Burning and stinging sensation, Eye irritation and infection, Eye pain, Post-injection flare, Wound healing suppression, Corneal perforation, Headache, Vitreous detachment, Subcapsular cataract, Ocular hypertension, Optic nerve damage, Visual field defect. Bacterial keratitis with the use of multiple dose containers.

Potential Contraindications / Warnings and Precautions

Infants and patients less than 12 years of age, Lactation, Pregnancy. Known hypersensitivity to any of components. Epithelial herpes simplex keratitis, Acute infectious stages of vaccinia, varicella and many other viral diseases of the corneal and conjunctiva. Mycobacterial infection of the eye, fungal diseases of the eye. Monitor IOP. Prolonged use may result in ocular hypertension and/or glaucoma and increase risk of secondary ocular infections.

Moxifloxacin HCl

Possible Adverse Reaction

Conjunctivitis, Decreased visual acuity, Dry eye, Keratitis, Ocular discomfort, Ocular hyperemia, Ocular pain, Ocular pruritus, Subconjunctival hemorrhage, and Tearing, Anaphylaxis, Growth of resistant organisms, Fever, Increased cough, Infection, Otitis media, Pharyngitis, Rash and Rhinitis.

Potential Contraindications / Warnings and Precautions

Patients less than 1 year, Lactation, Pregnancy. Known hypersensitivity to fluoroquinolones or any of the components. Growth of resistant organisms with prolonged use.