



Compound Formulation Active Pharmaceutical Ingredients and the Associated Adverse Effects and Potential Contraindications/Warnings and Precautions

Atropine Sulfate Monohydrate

Atropine Sulfate Monohydrate

Possible Adverse Reaction

Eye pain and stinging on administration, Blurred vision, Photophobia, Decreased lacrimation, superficial keratitis. Allergic reactions such as papillary conjunctivitis, contact dermatitis, and lid edema may also occur less commonly. Systemic side effects noted with Atropine 1%: Increased heart rate and blood pressure, dryness of skin, mouth, and throat from decreased secretions from mucus membranes; restlessness, irritability or delirium from stimulation of the central nervous system. Bacterial keratitis possible with multiple use container.

Potential Contraindications / Warnings and Precautions

Do not give to children less than 3 months of age. Patients 65 years of age or older, use this preparation with care. Pregnant or breast-feeding, talk to your physician prior to use. Potential for cross-sensitivity to any components. Use precaution if patient using MAOI (potential to precipitate hypertensive crisis).

*For professional use only. OSRX specializes in customizing compounded medications to meet unique patient needs. Compounded drugs are not FDA-approved, which means they have not undergone FDA premarket review for safety, effectiveness, and quality. **View potential adverse events and contraindications at: www.osrxpharmaceuticals.com/osrx-api-aecontraindication**

1120 Kensington Ave., Suite E, Missoula, MT 59801 | P: 855.466.1076 | F1: 406.541.6267 | F2: 866.515.0196 | osrxpharmaceuticals.com